

Nondecaying Hydrodynamic Interactions along Narrow Channels

Karolis Misiunas, Stefano Pagliara,^{*} Eric Lauga, John R. Lister, and Ulrich F. Keyser[†]
Cavendish Laboratory, University of Cambridge, United Kingdom
and Department of Applied Mathematics and Theoretical Physics, University of Cambridge,
Cambridge CB3 0HE, United Kingdom

(Received 16 February 2015; revised manuscript received 24 April 2015; published 15 July 2015)

Particle-particle interactions are of paramount importance in every multibody system as they determine the collective behavior and coupling strength. Many well-known interactions such as electrostatic, van der Waals, or screened Coulomb interactions, decay exponentially or with negative powers of the particle spacing r . Similarly, hydrodynamic interactions between particles undergoing Brownian motion decay as $1/r$ in bulk, and are assumed to decay in small channels. Such interactions are ubiquitous in biological and technological systems. Here we confine two particles undergoing Brownian motion in narrow, microfluidic channels and study their coupling through hydrodynamic interactions. Our experiments show that the hydrodynamic particle-particle interactions are distance independent in these channels. This finding is of fundamental importance for the interpretation of experiments where dense mixtures of particles or molecules diffuse through finite length, water-filled channels or pore networks.

DOI: 10.1103/PhysRevLett.115.038301

PACS numbers: 82.70.Dd, 47.57.eb, 47.60.-i, 83.50.Ha

Molecular diffusion inside channels and pores is relevant for a wide range of phenomena in natural systems [1] as well as technological applications [2,3]. For example, biological channels that transport antibiotics are under intense investigation due to their importance for drug transport [4]. Diffusing molecules inside protein channels are closely confined, leading to single-file diffusion with hydrodynamic interactions playing a role even at these nanometer length scales [5,6]. Transport through protein channels can be mimicked by colloidal particles and microfluidic chips [7] that confine particles to 1D diffusion [8]. Such particles undergo random walks in one dimension driven by Brownian motion, where the corresponding diffusion coefficient critically depends on the geometry of the confinement [3,9].

Loosely speaking, these Brownian particles receive momentum impulses from thermal fluctuations of the solvent molecules [10], and their resulting motion displaces the liquid around it [5,11]. This creates a flow field that mediates a long-range hydrodynamic interaction between the particles [12]. The strength of this interaction is proportional to the flow velocity which decays with distance as $1/r$, where r is the separation between two unconstrained particles in three dimensions (3D) [13]. Introducing a geometrical confinement modifies the flow field that changes the decay rate. For example, the interaction between particles constrained by two parallel plates (2D) decays faster at a rate of $\sim 1/r^2$ [14].

In narrow channels the hydrodynamic interactions were previously measured to rapidly decay with particle separation [15–17]. Indeed, the steady flow induced by particle motion in a channel decays exponentially with $r/2R$, where $2R$ is the channel width [5,11,18,19]. Consequently, the particle-particle interaction strength is expected to also decay exponentially with their separation [15,18]. However, recent theoretical investigation suggests that interactions could

have a longer spacial extent and a slower decay rate than previously thought [20]. Furthermore, previous experiments did not capture far field hydrodynamics because they used microfluidic chips with a groove geometry for mimicking channels [8,15–17]. Sedimentation kept the particles from escaping the microfluidic groove, but the liquid had no such constraint. The lack of controlled experiments in 1D confinement leaves many unanswered questions about the magnitude and spacial extent of hydrodynamic interactions inside narrow channels [8,21].

In this Letter, we present the first measurement of the interactions between two Brownian particles inside a finite narrow channel, that confines both the particles and the liquid. Figure 1 illustrates our experiments, where the “closed” channel, shown on the left, has only one end connected to a bulk reservoir, while the “open” channel, shown on the right, has both ends connected. Here we will demonstrate a fundamental difference between diffusion in open channels and in closed channels [11,15,18].

The experiments are realized using microfluidic lab-on-a-chip devices because they allow for direct imaging of particle motion. Inside the chip two large reservoirs are separated by a membrane containing closed [Fig. 2(a)] and open channels [Fig. 2(b)]. All channels are 5–17 μm long and have semielliptical cross sections with a height and width of approximately 800 nm that closely confine spherical particles of diameter $2a = 505 \pm 8$ nm. The resulting particle to channel size ratio is $a/R \approx 0.6$, ensuring particles always move in single file. Two additional large connections are positioned $\sim 200 \mu\text{m}$ away from the narrow channels allowing pressure equalization between the two reservoirs. For the fabrication of the chips, we use focused ion beam, photolithography, and replica molding of polydimethylsiloxane (PDMS) [22]. Crucially, the PDMS chips are then




FIG. 1 (color online). Trajectories of two particles undergoing Brownian motion in closed and open channels. The trajectories are visibly uncorrelated in the closed channel, suggesting the particles move independently. In the open channel the trajectories resemble each other, which leads to strong motion correlation.

oxygen plasma bonded onto a glass slide which provides a bottom wall confinement for the channels. Subsequently, we fill the chip with the polystyrene colloidal particles (Polysciences Inc.) dispersed in a 5 mM KCl salt solution that limits electrostatic interactions to a few nanometers.

An assembled chip is mounted onto an inverted, custom-built optical microscope with a high numerical aperture oil-immersion objective ($\times 100$; NA 1.4; UPLSAPO). Using holographic optical tweezers [23–25] we position two particles inside the channel, and then turn off the laser trapping which releases the particles and allows them to diffuse freely. Their motion is recorded using a CMOS camera (DMK-31BF03, Imaging Source) at a rate of 30 frames per second, until one particle escapes the channel. Afterwards, the particle trajectories are extracted from the images using standard image analysis techniques [26]. See Supplemental Material, videos S1 and S2 [27].

For data analysis, we divide the trajectories into displacement steps between consecutive frames: Δx_1 and Δx_2 denote the displacements of the first particle (one on the left) and the second particle (one on the right), respectively. The two-particle interaction strength is expected to be a function of their separation, and therefore, we group the pairs of displacements according to distance between the centers of the particles.

For the closed channel, Fig. 2(c) shows the displacements of the second particle (Δx_2) as a function of displacement of the first particle (Δx_1). The distribution is circular with points distributed equally in each quadrant, suggesting that the two particles move independently from each other. This is expected for a large particle separation of $10a$ [15,18]. In contrast, the distribution for the open channel, shown in Fig. 2(d), is elliptical with the major axis along $y = x$ and ellipticity of 0.75. This implies that the two particles move in the same direction more frequently than in opposite directions, suggesting the presence of long-ranged interaction between the particles.

We quantify the interaction strength using the Pearson product-moment correlation coefficient defined as [31]:




FIG. 2 (color online). Comparison between closed channel (left column) and open channel (right column) experiments. The top row (a),(b) shows bright field images of microfluidic channels containing two colloidal particles. Scale bars indicate $2 \mu\text{m}$. The middle row (c),(d) shows displacements of the second particle (\hat{y} axis) as a function of displacement of the first particle (\hat{x} axis). Each distribution contains 1000 displacement pairs with an initial particle separation of approximately $2.5 \mu\text{m}$. The overlaid lines indicate contours for σ and 2σ from a 2D normal-distribution fit. Insets illustrate the direction of motion for each quadrant. The bottom row (e),(f) shows the correlation coefficients versus the separation between the two particles. The solid lines indicate the fits to the phenomenological model. The two-particle motion is strongly correlated in the open channel, suggesting a presence of long-ranged nondecaying interaction.

$\rho = \text{cov}(\Delta x_1, \Delta x_2) / \sigma(\Delta x_1)\sigma(\Delta x_2)$, where cov is the covariance, and σ is the standard deviation. The value $\rho = 0$ indicates independent particle motion and $\rho = 1$ corresponds to fully correlated motion.

Figures 2(e) and 2(f) show the correlation coefficients as a function of particle separation. Evidently, the correlation is stronger and has a longer range in the open channel. A detailed examination of the closed channel results shows a high correlation when particles are close to one another that exponentially decays to zero in a separation of $\sim 4a$, as expected. Therefore, we fit a phenomenological model, $y = A \exp(-x/B) + C$, that captures the decay rate and has an additional offset parameter. The fit yields $B = 0.15 \pm .02 \mu\text{m}$, $C = 0.005 \pm .004$. We have introduced the offset to characterize the novel behavior observed in the open channel. As evident from the data, the correlation coefficient exhibits the same initial exponential decay, but in stark contrast, it asymptotes to a constant offset. This finite correlation coefficient is captured by the fit to the phenomenological model, yielding $B = 0.14 \pm .02 \mu\text{m}$,

$C = 0.419 \pm .005$. This constant, nondecaying component is not expected. Furthermore, the correlation persists even at the largest measured distances, suggesting that the two particles interact when both are inside the open channel. To the best of our knowledge, this is the first observation of such distance-independent interactions between Brownian particles.

We further investigate the effect of the channel length on the particle-particle interaction strength. Figure 3 shows the correlation coefficients for open channels of lengths $L = 5, 10, \text{ and } 17 \mu\text{m}$. The data clearly indicate that the interaction strength decreases with L . In the longest channel, shown as circles, we observe particles interacting at separations of more than $40a$. This is the largest relative distance measured between interacting Brownian particles, even exceeding the $\sim 20a$ separation measured in bulk [13,32]. This is a surprising result because the geometric confinement typically reduces the maximum interaction distance [11,15,32].

Based on our observations we propose a hypothesis for the distance-independent interaction mechanism, that is schematically illustrated in the inset of Fig. 3. Suppose the first particle, shown on the left, moves to the right due to a thermal momentum impulse. At steady state its motion induces a flow that is constrained by the channel geometry to flow either around the particle or along the channel. Importantly, the latter flow has been previously neglected [5,11,15,18] because the studies considered infinitely long channels, in which the finite pressure exerted by the moving particle cannot displace an infinite liquid column. In contrast, we argue that for finite open channels, flows extend throughout the whole channel. Consequently, the induced flow along the channel has a constant mean flow velocity that is proportional to the driving force, i.e., the first particle's velocity [11,33]. This flow encounters the second particle on the right, and exerts a viscous




FIG. 3 (color online). Correlation coefficients for two interacting particles for different channel lengths. The three curves correspond to open channels with different lengths, from top to bottom: 5, 10, 17 μm . The solid lines show the fit to a model $y = A \exp(-x/B) + C$, where the offset coefficients were (from top to bottom): $C = 0.420 \pm .006$; $0.250 \pm .005$; $0.101 \pm .004$. Evidently, the long-ranged two-particle correlation coefficient decreases with the channel length. Inset illustrates the proposed model for long-ranged hydrodynamic interaction.

drag force on it. The magnitude of this force is a function of flow velocity [34] and thus also a function of the first particle's velocity, but is independent of the particle positions. The opposite case of the second particle moving can be inferred by symmetry, and the final interaction is a combination of the two cases. This gives rise to the particle-particle interaction that we observed with our correlation coefficient measurements and also explains why the value is distance-independent in open channels. In a limit where the particle size matches the channel, this problem resembles two pistons in a pipe and the trajectories would be perfectly correlated [12]. On the contrary, in the closed channels, the dead end blocks the flow along the channel, thus eliminating the nondecaying hydrodynamic interaction.

We now use this hypothesis to construct an analytical model that sheds light on the physics of the interaction. The model focuses on physical scalings and omits detailed numerical prefactors from the presentation. Our goal is to estimate the typical mean flow velocity in the channel resulting from the motion of one particle.

Consider a spherical particle of diameter $2a$ located in the center of an open cylindrical channel of radius R and length L . We assume that the particle moves instantaneously to the right with velocity U . The fluid displaced by the particle must either be pushed along the channel to the right channel end, with more fluid drawn in at the left, or leak from right to left through the thin gap between the particle and the channel wall. The pressure increase across the sphere, Δp , is proportional to the flow rate in the channel, Q , according to Poiseuille law

$$\Delta p \sim \frac{\mu Q L}{R^4}, \quad (1)$$

where μ is the dynamic viscosity. We neglect any hydrodynamic resistance due to the recirculation from the exit of the channel to the entrance, equivalent to imposing a periodic boundary condition. Mass conservation around the moving sphere (in the frame moving with the sphere) leads to

$$Q - UR^2 \sim Rq, \quad (2)$$

where q is the leakage flux through the thin gap between the sphere and the channel. It is approximately given by lubrication theory [35]

$$q \sim -Uh - \frac{\Delta p h^3}{\mu l_0}, \quad (3)$$

where $h \equiv R - a$ is the minimum gap width, and $l_0 \sim (ah)^{1/2}$ is the characteristic lubrication length scale. Combining these equations and taking the limit $h \ll R$, we obtain the flow rate

$$Q \sim UR^2 \left(\frac{R^3 a^{1/2}}{R^3 a^{1/2} + L h^{5/2}} \right). \quad (4)$$

Combining with the mean flow in the channel, $\langle u \rangle \sim Q/R^2$, we get the final expression

$$\frac{\langle u \rangle}{U} \sim \frac{R^3 a}{R^3 a + L h^{5/2} a^{1/2}}. \quad (5)$$

Our result shows that the mean flow in the channel is nonzero, but a function of the channel dimensions R and L . Importantly, our model reproduces the observed decrease with the channel length: scaling with $1/L$. The long channel limit ($L \rightarrow \infty$) gives no external flows ($Q = 0$) equivalent to the closed channel case.

In order to quantitatively capture the experimental results, we employ numerical simulations. Using finite element analysis software (COMSOL MULTIPHYSICS v4.4) [9] we solve the Stokes equations for two spherical particles inside a cylindrical channel with a diameter chosen such that the areas match the experimental parameters. Periodic boundary conditions are set on the ends while all other surfaces were set to no-slip boundary conditions. We apply an instantaneous velocity on one particle and calculate the resulting drag forces on both particles. A velocity independent measure of these forces is known as a friction matrix (ζ) ([33], p. 226). For the two particle system in one dimension it is a 2×2 symmetric matrix, where the diagonal terms ($\zeta_{1,1}; \zeta_{2,2}$) describe hydrodynamic drag experienced by the moving particles (first; second), and the off-diagonal elements ($\zeta_{1,2} = \zeta_{2,1}$) correspond to a force exerted by the moving particle on the other particle. The friction matrix values are computed directly from the numerical simulation. In addition, the friction matrix can be obtained from the experiments using a diffusivity matrix, $\mathbf{D} = k_B T \zeta^{-1}$ [33], with components $D_{i,i} = \langle \Delta x_i^2 \rangle / 2\Delta t$ and $D_{1,2} = D_{2,1} = \langle \Delta x_1 \Delta x_2 \rangle / 2\Delta t$ [36]. This allows us to compare the hydrodynamic interactions predicted by our simulation with the measured data.

Figures 4(a) and 4(b) illustrate the typical flow patterns computed with our simulation. Notice that the flow in the closed channel curls around the particle and does not extend far into the channel. This contrasts with the open channel, where the flow around the particle is weaker and there is a Poiseuille flow along the whole channel length. This moving fluid column exerts a force on the second particle that gives rise to the nondecaying interaction. Figures 4(c) and 4(d) show a quantitative comparison between the numerical simulation and experimental data, where the solid lines indicate the simulation results and the points indicate the experimental values. The friction with the channel walls is the same for both particles, leading to the overlapping curves ($\zeta_{1,1}$ and $\zeta_{2,2}$). Meanwhile,

the interaction force ($\zeta_{1,2}$) asymptotes to a nonzero value for the open channel only, similar to the correlation coefficient that was reported above. At small separations the discrepancy between simulation and experiment is likely caused by electrostatics and finite frame rate. Meanwhile, at large separations the values compare very well, with the largest discrepancy below 20%. This is very good agreement given the approximations made about channel shape and width.

The agreement between the numerical simulation and the experimental data suggest that the proposed hypothesis captures the physics of nondecaying interactions between the Brownian particles. Crucially, this interaction requires flow through the ends of the channel, which is attained by keeping the two reservoirs at equal pressures. In our experiments, the pressure equalizes via liquid recirculation through the secondary large channels placed far away from the narrow channel. In biological cells, this can happen through the membrane or through dedicated water transport protein channels known as aquaporins [1]. In porous materials, the pressure can equalize through other interlinking channels [2].

Our observations have far-reaching implications for diffusion processes inside channels. One of the predictions from our model is that the viscous drag on particles is smaller than the theory expects for narrow channels (Fig. S5). We therefore performed an additional experiment that measured the diffusion coefficient of a single particle inside an $L = 10 \mu\text{m}$ open channel (method [9]). The measured diffusion coefficient is indeed 40% higher than expected [5,11]: $D_x/D_0 = 0.126 \pm .006$, where D_0 is the diffusion coefficient in the bulk (Fig. S6). This potentially explains the experimental discrepancies noted in previous studies for channels with similar dimensions [9,37,38]. Because the diffusion coefficient is important for predicting transport rates, our results directly impact channel transport models [39]. Moreover, the long-ranged interaction between particles should lead to a cooperative behavior that could enhance transport across channels. Also, since the interactions are not limited to two particles, they should persist for channels filled with three or even more particles. This makes our results relevant for polymers in confinement because the monomers can interact nonlocally [40],




FIG. 4 (color online). Top row illustrates flows inside the (a) closed and (b) open channel obtained from the simulations. The bottom row compares the friction matrix values for the (c) closed and (d) open channel. The points indicate experimentally measured values, while solid lines show the corresponding simulation results obtained without any fitting parameters. Values are normalized with the bulk coefficient: $\zeta_0 = 6\pi\mu a$. Arrows highlight the non-zero interaction term which agrees well with our simulation results.

and also for single-file systems because the particles interact not only with the closest neighbors [8].

In conclusion, we utilized a highly controlled microfluidic system coupled with holographic optical tweezers to investigate the interaction of particles in confinement. Our measurements prove that interactions extend over the full channel length and have a constant strength that does not decay with particle separation. We explain the coupling mechanism using hydrodynamics with both a qualitative analytical model and quantitative comparisons with the numerical simulations. The excellent agreement between the theory and experiments suggests that we fully understand the properties of hydrodynamic particle interactions in microfluidic channels. The nondecaying interaction extending throughout the whole channel has important implications for the modeling of transport through channels as well as for the interpretation of experiments investigating particles diffusing in close confinement.

All data accompanying this publication are directly available within the publication.

The authors thank S. Ghosal and M. Muthukumar for fruitful discussions. U. F. K. was supported by an ERC starting Grant No. (PassMembrane 261101). S. P. acknowledges funding from a Leverhulme Early Career Fellowship. K. M. was supported by a grant from the EPSRC. E. L. was supported by a Marie Curie CIG grant from the EU.

*Present address: Department of Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom.

†To whom all correspondence should be addressed.
ufk20@cam.ac.uk

- [1] B. Alberts, A. Johnson, J. Lewis, D. Morgan, M. Raff, K. Roberts, and P. Walter, *Molecular Biology of the Cell*, 6th ed. (Garland Science, New York, 2014).
- [2] J. Kärger, T. Binder, C. Chmelik, F. Hibbe, H. Krautscheid, R. Krishna, and J. Weitkamp, Microimaging of transient guest proles to monitor mass transfer in nanoporous materials, *Nat. Mater.* **13**, 333 (2014).
- [3] T. Motz, G. Schmid, P. Hänggi, D. Reguera, and J. M. Rubí, Optimizing the performance of the entropic splitter for particle separation, *J. Chem. Phys.* **141**, 074104 (2014).
- [4] J.-M. Pagès, C. E. James, and M. Winterhalter, The porin and the permeating antibiotic: a selective diusion barrier in Gram-negative bacteria, *Nat. Rev. Microbiol.* **6**, 893 (2008).
- [5] W. Deen, Hindered transport of large molecules in liquidfilled pores, *AIChE J.* **33**, 1409 (1987).
- [6] S. Gravelle, L. Joly, F. Detcheverry, C. Ybert, C. Cottin-Bizonne, and L. Bocquet, Optimizing water permeability through the hourglass shape of aquaporins, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 16367 (2013).
- [7] S. Pagliara, S. L. Dettmer, and U. F. Keyser, Channel-Facilitated Diusion Boosted by Particle Binding at the Channel Entrance, *Phys. Rev. Lett.* **113**, 048102 (2014).
- [8] Q. Wei, C. Bechinger, and P. Leiderer, Single-file diusion of colloids in one-dimensional channels, *Science* **287**, 625 (2000).
- [9] S. L. Dettmer, S. Pagliara, K. Misiunas, and U. F. Keyser, Anisotropic diusion of spherical particles in closely conning microchannels, *Phys. Rev. E* **89**, 062305 (2014).
- [10] R. Huang, I. Chavez, K. M. Taute, B. Lukic, S. Jeney, M. G. Raizen, and E.-L. Florin, Direct observation of ballistic Brownian motion on a single particle, *Nat. Phys.* **7**, 576 (2011).
- [11] J. Happel and H. Brenner, *Low Reynolds Number Hydrodynamics*, edited by R. Moreau (Martinus Nijhoff Publishers, The Hague, 1973).
- [12] H. Diamant, Hydrodynamic interaction in conned geometries, *J. Phys. Soc. Jpn.* **78**, 041002 (2009).
- [13] J. C. Crocker, Measurement of the hydrodynamic corrections to the Brownian motion of two colloidal spheres, *J. Chem. Phys.* **106**, 2837 (1997).
- [14] B. Cui, H. Diamant, B. Lin, and S. A. Rice, Anomalous Hydrodynamic Interaction in a Quasi-Two-Dimensional Suspension, *Phys. Rev. Lett.* **92**, 258301 (2004).
- [15] B. Cui, H. Diamant, and B. Lin, Screened Hydrodynamic Interaction in a Narrow Channel, *Phys. Rev. Lett.* **89**, 188302 (2002).
- [16] D. T. Valley, S. a. Rice, B. Cui, H. M. Ho, H. Diamant, and B. Lin, Pair diusion in quasi-one- and quasi-two-dimensional binary colloid suspensions, *J. Chem. Phys.* **126**, 134908 (2007).
- [17] E. Kosheleva, B. Leahy, H. Diamant, B. Lin, and S. A. Rice, Long-range hydrodynamic correlations in quasi-one-dimensional circular and straight geometries, *Phys. Rev. E* **86**, 041402 (2012).
- [18] N. Liron and R. Shahar, Stokes ow due to a Stokeslet in a pipe, *J. Fluid Mech.* **86**, 727 (1978).
- [19] N. Al Quddus, W. a. Moussa, and S. Bhattacharjee, Motion of a spherical particle in a cylindrical channel using arbitrary Lagrangian-Eulerian method, *J. Colloid Interface Sci.* **317**, 620 (2008).
- [20] D. Frydel and H. Diamant, Long-Range Dynamic Correlations in Conned Suspensions, *Phys. Rev. Lett.* **104**, 248302 (2010).
- [21] D. Reguera, G. Schmid, P. S. Burada, J. M. Rubí, P. Reimann, and P. Hänggi, Entropic Transport: Kinetics, Scaling, and Control Mechanisms, *Phys. Rev. Lett.* **96**, 130603 (2006).
- [22] S. Pagliara, C. Chimere, R. Langford, D. G. a. L. Aarts, and U. F. Keyser, Parallel sub-micrometre channels with dierent dimensions for laser scattering detection, *Lab Chip* **11**, 3365 (2011).
- [23] D. G. Grier, A revolution in optical manipulation, *Nature (London)* **424**, 810 (2003).
- [24] M. Padgett and R. Di Leonardo, Holographic optical tweezers and their relevance to lab on chip devices, *Lab Chip* **11**, 1196 (2011).
- [25] S. Pagliara, C. Schwall, and U. F. Keyser, Optimizing diusive transport through a synthetic membrane channel, *Adv. Mater.* **25**, 844 (2013).
- [26] S. L. Dettmer, U. F. Keyser, and S. Pagliara, Local characterization of hindered Brownian motion by using digital video microscopy and 3D particle tracking, *Rev. Sci. Instrum.* **85**, 023708 (2014).
- [27] See Supplemental Material <http://link.aps.org/supplemental/10.1103/PhysRevLett.115.038301> for video samples of our experiments, supporting experiments, and details on our simulations, which includes Refs. [28–30].

- [28] H. Brenner, Macrotransport processes, *Langmuir* **6**, 1715 (1990).
- [29] D. L. Ermak and J. A. McCammon, Brownian dynamics with hydrodynamic interactions, *J. Chem. Phys.* **69**, 1352 (1978).
- [30] T. Savin and P. S. Doyle, Static and dynamic errors in particle tracking microrheology, *Biophys. J.* **88**, 623 (2005).
- [31] K. F. Riley, P. Hobson, and S. J. Bence, *Mathematical Methods for Physics and Engineering: A Comprehensive Guide* (Cambridge University Press, Cambridge, England, 2006).
- [32] E. R. Dufresne, T. M. Squires, M. P. Brenner, and D. G. Grier, Hydrodynamic Coupling of Two Brownian Spheres to a Planar Surface, *Phys. Rev. Lett.* **85**, 3317 (2000).
- [33] R. M. Mazo, *Brownian Motion: Fluctuations, Dynamics, and Applications* (Clarendon, Oxford University Press, Oxford, 2002).
- [34] W. R. Smythe, Flow around a sphere in a circular tube, *Phys. Fluids* **4**, 756 (1961).
- [35] G. K. Batchelor, *An Introduction to Fluid Dynamics*, *Cambridge Mathematical Library* (Cambridge University Press, Cambridge, England, 2000).
- [36] X. Xu, S. A. Rice, B. Lin, and H. Diamant, Influence of Hydrodynamic Coupling on Pair Diffusion in a Quasi-One-Dimensional Colloid System, *Phys. Rev. Lett.* **95**, 158301 (2005).
- [37] S. Pagliara, S. L. Dettmer, K. Misiunas, L. Lea, Y. Tan, and U. F. Keyser, Diffusion coefficients and particle transport in synthetic membrane channels, *Eur. Phys. J. Spec. Top.* **223**, 3145 (2014).
- [38] M. Schiel and Z. S. Siwy, Diffusion and trapping of single particles in pores with combined pressure and dynamic voltage, *J. Phys. Chem. C* **118**, 19214 (2014).
- [39] A. M. Berezhkovskii and S. M. Bezrukov, Optimizing transport of metabolites through large channels: molecular sieves with and without binding, *Biophys. J.* **88**, L17 (2005).
- [40] P.-G. de Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca, NY, 1979).